arXiv:1704.06015v1 [gr-qc] 20 Apr 2017

PEELING PROPERTY OF BONDI-SACHS METRICS FOR NONZERO COSMOLOGICAL CONSTANT

FANGQUAN XIE, XIAO ZHANG

Abstract. In this short paper, we show that the peeling property still holds for Bondi-Sachs metrics with nonzero cosmological constant under the new boundary condition with nontrivial B, X, Y obtained in [6]. This should indicate the new boundary condition is natural. Moreover, we construct some nontrivial vacuum Bondi-Sachs metrics without the Bondi news. This provides a new feature of gravitational waves for nonzero cosmological constant.

PACS(2010): 04.70Bw, 04.25Dm, 04.20Cv

1. Introduction

The full nonlinear theory of gravitational waves was established until the work of Bondi, Sachs, et al. in framework of Bondi-Sachs metrics [1, 2]. Lately, the theory was re-formulated by Penrose in a more geometric notion of conformal compactification of spacetimes [3]. However, they were well-studied only when the cosmological constant $\Lambda = 0$. In this case, by assuming Sommerfeld's radiation condition, there is a natural boundary condition to make the vacuum Bondi-Sachs metrics asymptotic to

$$\begin{aligned} g =& \Big(1 - \frac{2M}{r}\Big) du^2 + 2dudr - 2ldud\theta - 2\bar{l}\sin\theta dud\psi \\ & - r^2\Big[(1 + \frac{2c}{r})d\theta^2 + \frac{4d}{r}\sin\theta d\theta d\psi + (1 - \frac{2c}{r})\sin^2\theta d\psi^2\Big] \end{aligned}$$

where u is retarded coordinate, r, θ and ψ are polar coordinates, M, c, d are smooth functions of u, θ, ψ defined on $\mathbb{R} \times S^2$ with regularity condition $\int_0^{2\pi} c(u, \theta, \psi) d\psi = 0$ for $\theta = 0, \pi$ and for all u, $l = c_{,\theta} + 2c\cot\theta + d_{,\psi}\csc\theta$, $\bar{l} = d_{,\theta} + 2d\cot\theta - c_{,\psi}\csc\theta$.

The functions c, d are the *Bondi news* which characterizes the presence of gravitational waves. Geometrically, they measure the deformation of standard 2-sphere as a result of gravitational waves. The Bondi energy-momentum defined on u-slice is probably the most important feature of the nonlinear theory

$$m_\nu(u) = \frac{1}{4\pi}\int_{S^2} M(u, \theta, \psi) n^\nu dS, \qquad \nu = 0, 1, 2, 3,$$

where $n^0 = 1$, $n^1 = \sin\theta\cos\psi$, $n^2 = \sin\theta\sin\psi$, $n^3 = \cos\theta$. The famous Bondi energy loss formula asserts [1, 2]

$$\frac{d}{du}m_0(u) = -\frac{1}{4\pi}\int_{S^2}(c_{,u})^2 + (d_{,u})^2 \leq 0.$$

Indeed, if $|m(u)| = \sqrt{m_1^2(u) + m_2^2(u) + m_3^2(u)} \neq 0$, then it was improved to the Bondi energy-momentum loss [4, 5]

$$\frac{d}{du}\Big(m_0(u) - |m(u)|\Big) = -\frac{1}{4\pi}\int_{S^2}\Big[(c_{,u})^2 + (d_{,u})^2\Big]\Big(1 - \frac{m_i n^i}{|m|}\Big) \leq 0.$$

Therefore the Bondi energy-momentum can be viewed as the total energy-momentum measured after the loss due to the gravitational radiation up to that time.

It was shown, under this natural boundary condition, the Weyl curvature components of Bondi-Sachs metrics satisfy the *peeling* property

$$\Psi_k = O\Big(\frac{1}{r^{5-k}}\Big)$$

in the case $\Lambda = 0$. This property is crucial for constructing wave forms in numerical simulations.

As cosmological observations indicated that the universe has a positive cosmological constant, it has been studied extensively for the nonlinear theory of gravitational waves in recent years, e.g. [6, 7, 8, 9, 10, 11]. In [6], a detail asymptotic analysis of Bondi-Sachs metrics was provided. By assuming again Sommerfeld's radiation condition, which is natural in numerical simulations, the new boundary condition with three additional functions B, X, Y were discussed and, in particular, the nonzero Bondi news causes X, Y to be nonzero. In [7], an alternative boundary condition was provided in the axi-symmetric case, without assuming Sommerfeld's radiation condition, but deforming 2-sphere with certain factors involving in cosmological constant. In a series of papers [8, 9, 10, 11], asymptotics with a positive cosmological constant was discussed in framework of Penrose, and the linearization theory as well as the quadrupole formula were also derived. Some revelent works on the linearization theory can also be found in [12, 13, 14].

In this short paper, we show that the peeling property still holds for Bondi-Sachs metrics with nonzero cosmological constant under the new boundary condition with nontrivial B, X, Y obtained in [6]. It is somehow surprising as the Bondi-Sachs metrics do not seem to be (anti-)de Sitter at infinity in current situation. But the peeling property should indicate the new boundary condition is natural. Moreover, we construct some nontrivial vacuum Bondi-Sachs metrics without the Bondi news. This provides a new feature of gravitational waves for nonzero cosmological constant, and it implies some class of gravitational waves may be missed in detection by measuring the Bondi news in the case of positive cosmological constant.

2. Natural boundary condition

Suppose a spacetime has a family of non-intersecting null hypersurfaces given by the level sets of smooth function u. We choose coordinates $x^0 = u$, $x^1 = r$, $x^2 = \theta$ and $x^3 = \phi$ where r is a luminosity distance along the null rays. Following from [6], Bondi-Sachs metrics [1, 2], even with the nonzero cosmological constant Λ, are

$$g = \Big(e^{2\beta}\frac{V}{r} - r^2 h_{AB}U^A U^B\Big)du^2 + 2e^{2\beta}dudr + 2r^2 h_{AB}U^B du dx^A - r^2 h_{AB}dx^A dx^B$$

where $A, B = 2, 3$, $U^2 = U$, $U^3 = W\csc\theta$,

$$h \equiv \begin{pmatrix} h_{22} & h_{23} \\ h_{32} & h_{33} \end{pmatrix} \equiv \begin{pmatrix} e^{2\gamma}\cosh 2\delta & \sinh 2\delta \sin\theta \\ \sinh 2\delta\sin\theta & e^{-2\gamma}\cosh 2\delta \sin^2\theta \end{pmatrix},$$

β, γ, δ, V, U, W are functions of u, r and points on unit 2-sphere parameterized by θ, ϕ, i.e., they take the same values at $\phi = 0$ and $\phi = 2\pi$.

Now we study the vacuum Einstein field equations

$$R_{ij} + \Lambda g_{ij} = 0$$

If we assume Sommerfeld's radiation condition, i.e., γ, δ can be expanded as follows

$$\gamma = \frac{c}{r} + \Big(-\frac{1}{6}c^3 - \frac{3}{2}d^2 c + C\Big)\frac{1}{r^3} + O\Big(\frac{1}{r^4}\Big),$$
$$\delta = \frac{d}{r} + \Big(-\frac{1}{6}d^3 + \frac{1}{2}c^2 d + D\Big)\frac{1}{r^3} + O\Big(\frac{1}{r^4}\Big)$$

with the regularity condition $\int_0^{2\pi} c(u,\theta,\psi)d\psi = 0$ for $\theta = 0, \pi$ and for all u, then we can obtain the following asymptotic expansions [6],

$$\begin{aligned}
\beta =& B - \frac{c^2 + d^2}{4r^2} + O\Big(\frac{1}{r^3}\Big) \\
W =& X + 2e^{2B}B_{,\phi}\csc\theta\frac{1}{r} + e^{2B}\Big(2cB_{,\phi}\csc\theta - 2dB_{,\theta} - \bar{l}\Big)\frac{1}{r^2} + O\Big(\frac{1}{r^3}\Big) \\
U =& Y + 2e^{2B}B_{,\theta}\frac{1}{r} - e^{2B}\Big(2dB_{,\phi}\csc\theta + 2cB_{,\theta} + l\Big)\frac{1}{r^2} + O\Big(\frac{1}{r^3}\Big) \\
V =& -\frac{e^{2B}\Lambda}{3}r^3 + (\cot\theta Y + \csc\theta X_{,\phi} + Y_{,\theta})r^2 + e^{2B}\big(4B_{,\phi}^2\csc^2\theta + 2B_{,\phi\phi}\csc^2\theta \\
& + 2B_{,\theta}\cot\theta + 4B_{,\theta}^2 + 2B_{,\theta\theta} + 1\big)r - 2M + O\Big(\frac{1}{r}\Big)
\end{aligned}$$

where $l = c_{,\theta} + 2c\cot\theta + d_{,\psi}\csc\theta$, $\bar{l} = d_{,\theta} + 2d\cot\theta - c_{,\psi}\csc\theta$. Moreover, X, Y satisfy

$$\csc\theta X_{,\phi} + \cot\theta Y - Y_{,\theta} = -\frac{2}{3}e^{2B}\Lambda c, \quad X_{,\theta} - \cot\theta X + \csc\theta Y_{,\phi} = \frac{2}{3}e^{2B}\Lambda d.$$

If the cosmological constant $\Lambda = 0$, then X, Y must be zero, and B can be chosen to zero by suitable coordinate transformation [1, 2]. However, when $\Lambda \neq 0$, the situation is completely different. If the Bondi news c, d are nonzero, they will cause

X, Y to be nonzero according to the above equations. Moreover, B can not be transformed to zero in general. This provides a new boundary condition. In this case, we let $X = \Lambda a \sin\theta$, $Y = \Lambda b \sin\theta$, the above equations give

$$a_{,\phi} - \sin\theta b_{,\theta} = -\frac{2}{3}e^{2B}c, \qquad b_{,\phi} + \sin\theta a_{,\theta} = \frac{2}{3}e^{2B}d$$

and a, b can be determined uniquely by B, c, d up to some functions $\rho(u)$, $\sigma(u)$.

3. Peeling property

In this section we show the peeling property still holds for the new boundary condition when the cosmological constant is nonzero. This is somehow surprising as the new boundary condition does not indicate the Bondi-Sachs metrics are de-Sitter or anti-de Sitter at infinity.

Choose the null tetrad of Bondi-Sachs metrics

$$\begin{aligned}
e_0 &= \mathbf{l} = e^{-2\beta}\Big(\frac{\partial}{\partial u} - \frac{V}{2r}\frac{\partial}{\partial r} + U\frac{\partial}{\partial\theta} + \csc\theta W\frac{\partial}{\partial\phi}\Big)\\
e_1 &= \mathbf{k} = \frac{\partial}{\partial r}\\
e_2 &= \mathbf{m} = \frac{e^{-\gamma}}{r\sqrt{2\cosh 2\delta}}\Big(1 - i\sinh 2\delta\Big)\frac{\partial}{\partial\theta} + i\frac{e^{\gamma}\sqrt{\cosh 2\delta}\csc\theta}{\sqrt{2}r}\frac{\partial}{\partial\phi}\\
e_3 &= \bar{\mathbf{m}} = \frac{e^{-\gamma}}{r\sqrt{2\cosh 2\delta}}\Big(1 + i\sinh 2\delta\Big)\frac{\partial}{\partial\theta} - i\frac{e^{\gamma}\sqrt{\cosh 2\delta}\csc\theta}{\sqrt{2}r}\frac{\partial}{\partial\phi}
\end{aligned}$$

with the dual tetrad ω^0, ω^1, ω^2, ω^3. Then the metric is

$$ds^2 = 2\omega^0\omega^1 - 2\omega^2\omega^3.$$

The structure coefficients of the tetrad is denoted by C_{ij}^k which satisfy the commutation conditions

$$[e_i, e_j] = C_{ij}^k e_k.$$

By the basic formula of Riemannian geometry

$$\begin{aligned}
2\langle\nabla_X Y, Z\rangle =& X\langle Y, Z\rangle + Y\langle Z, X\rangle - Z\langle X, Y\rangle\\
=& \langle[X, Y], Z\rangle - \langle[Y, Z], X\rangle + \langle[Z, X], Y\rangle,
\end{aligned}$$

the connection coefficients and the structure coefficients satisfy

$$\Gamma_{ijk} = \frac{1}{2}\big(C_{ijk} - C_{jki} + C_{kij}\big)$$

where $\Gamma_{ijk} = \Gamma_{ij}^h g_{hk}$, $C_{ijk} = C_{ij}^h g_{hk}$. The Riemann curvature tensor satisfies

$$R_{jkl}^i = e_k(\Gamma_{lj}^i) + \Gamma_{lj}^h\Gamma_{kh}^i - C_{kl}^h\Gamma_{hj}^i - e_l(\Gamma_{kj}^i) - \Gamma_{kj}^h\Gamma_{lh}^i.$$

The vacuum Einstein field equations show that the Weyl tensor $\mathcal{C}_{ijkl}$ satisfies

$$\mathcal{C}_{ijkl} = R_{ijkl} + \frac{2}{3}\Lambda g_{i[k}g_{l]j}.$$

The Newmann-Penrose quantity Φ_k, $k = 0, \dots, 4$, are defined by

$$\begin{aligned}
\Psi_0 &= \mathcal{C}_{1212} = R_{1212},\\
\Psi_1 &= \mathcal{C}_{1012} = R_{1012},\\
\Psi_2 &= \mathcal{C}_{1230} = R_{1230} + \frac{\Lambda}{3},\\
\Psi_3 &= \mathcal{C}_{0103} = R_{0103},\\
\Psi_4 &= \mathcal{C}_{0303} = R_{0303}.
\end{aligned}$$

Peeling Property: *Under the boundary condition with nonzero B, X and Y,*

$$\Psi_k = \Big[\big(\Psi_k^{5-k}\big)^0 + \big(\Psi_k^{5-k}\big)^1\Lambda + O\big(\Lambda^2\big)\Big]\frac{1}{r^{5-k}} + O\Big(\frac{1}{r^{6-k}}\Big)$$

where coefficients $\big(\Psi_k^{5-k}\big)^0$, $\big(\Psi_k^{5-k}\big)^1$, ..., are given by B, a, b and other functions appeared in the expansions of Bondi-Sachs metrics, together with their derivatives. These coefficients have the same physical scale.

The peeling property can be proved by a direct computation. It is mysterious that un-falloff leading term of Ψ_4 and slow falloff leading terms of other Ψ_i vanish when $c = -\frac{3}{2}e^{-2B}(a_{,\phi} - \sin\theta b_{,\theta})$, $d = \frac{3}{2}e^{-2B}(b_{,\phi} + \sin\theta a_{,\theta})$ are substituted. The precise terms of $\big(\Psi_k^{5-k}\big)^0$ are given in the appendix.

4. Gravitational waves without the Bondi news

In this section we construct some nontrivial vacuum Bondi-Sachs metrics with $\gamma = \delta = 0$. Hence the Bondi news is zero. As in [6], taking the axi-symmetric Bondi-Sachs metrics with $X = 0$, $Y = \sin\theta\sigma(u)$, $M = \tau(u)e^{-4B}$, then the metrics

$$\begin{aligned}
g =& e^{2B}\Big[-\frac{\Lambda}{3}e^{2B}r^2 + 2\cos\theta\sigma r + e^{2B}(4B_{,\theta}^2 + 2\cot\theta B_{,\theta} + 2B_{,\theta\theta} + 1) - \frac{2M}{r}\Big]du^2\\
&+ 2e^{2B}dudr - r^2\Big[(\sin\theta\sigma + \frac{2e^{2B}B_{,\theta}}{r})du - d\theta\Big]^2 - r^2\sin^2\theta d\phi^2
\end{aligned} \tag{4.1}$$

are vacuum if and only if $B(u,\theta)$ satisfies

$$\begin{aligned}
&e^{4B}\Big[16\cot\theta B_{,\theta}^3 + 4B_{,\theta\theta}^2 - 4B_{,\theta}^2(\cot^2\theta - 4B_{,\theta\theta}) - B_{,\theta\theta}\cot^2\theta\\
&+ 2\cot\theta B_{,\theta\theta\theta} + B_{,\theta}\big(\cot\theta(\csc^2\theta + 16B_{,\theta\theta} + 2) + 8B_{,\theta\theta\theta}\big) + B_{,\theta\theta\theta\theta}\Big]\\
&+ 4MB_{,u} - 2M_{,u} - 6\cos\theta M\sigma - 3\sin\theta M_{,\theta}\sigma = 0.
\end{aligned} \tag{4.2}$$

For any function $|C(u)| > 1$, we choose

$$B = \frac{\ln(C(u) + \cos\theta)}{2}, \quad M = \frac{\tau(u)}{(C(u) + \cos\theta)^2}.$$

It is easy to check that this B satisfies the following equation

$$\begin{aligned}&16\cot\theta B_{,\theta}^3+4B_{,\theta\theta}^2-4B_{,\theta}^2(\cot^2\theta-4B_{,\theta\theta})-B_{,\theta\theta}\cot^2\theta\\&+2\cot\theta B_{,\theta\theta\theta}+B_{,\theta}\big(\cot\theta(\csc^2\theta+16B_{,\theta\theta}+2)+8B_{,\theta\theta\theta}\big)+B_{,\theta\theta\theta\theta}=0.\end{aligned}$$

Now we derive σ so that the equation (4.2) holds. For the above B and M,

$$B_u=\frac{C'}{2(C+\cos\theta)},\ M_u=\frac{\tau'}{(C+\cos\theta)^2}-\frac{2\tau C'}{(C+\cos\theta)^3},\ M_\theta=\frac{2\tau\sin\theta}{(C+\cos\theta)^3}.$$

Thus (4.2) reduces to

$$\frac{2\tau C'}{(C+\cos\theta)^3}-\frac{2\tau'}{(C+\cos\theta)^2}+\frac{4\tau C'}{(C+\cos\theta)^3}-\frac{6\tau\sigma\cos\theta}{(C+\cos\theta)^2}-\frac{6\tau\sigma\sin^2\theta}{(C+\cos\theta)^3}=0.$$

Therefore

$$3\tau C'-\tau' C-3\tau\sigma-\big(\tau'+3\tau\sigma C\big)\cos\theta=0.$$

As τ, σ, C are functions of u, it gives

$$3\tau C'-\tau' C-3\tau\sigma=0,\quad \tau'+3\tau\sigma C=0.$$

Thus

$$\sigma=-\frac{C'}{C^2-1},\quad \tau=m\big(C^2-1\big)^{\frac{3}{2}}$$

where m is constant.

Substituting them into (4.1), one gets the following regular, exact vacuum Bondi-Sachs metrics without the Bondi news

$$\begin{aligned}g=&\Big[-(C+\cos\theta)^2\frac{\Lambda}{3}r^2-\frac{2\cos\theta(C+\cos\theta)C'}{C^2-1}+C^2-\cos^2\theta\\&-\frac{2m(C^2-1)^{\frac{3}{2}}}{r(C+\cos\theta)}-\sin^2\theta\Big(\frac{1-C^2-rC'}{C^2-1}\Big)^2\Big]du^2+2\big(C+\cos\theta\big)dudr\\&+2r\sin\theta\frac{1-C^2-rC'}{C^2-1}dud\theta-r^2\Big(d\theta^2+\sin^2\theta d\phi^2\Big).\end{aligned}$$

5. Appendix

The formulas of $\big(\Psi_k^{5-k}\big)^0$ are as follows.

$$
\begin{aligned}
(\Psi_4^1)^0 =& e^{-6B}\Big(-\frac{3}{2}a_{uu\phi} + \frac{3}{2}b_{uu\phi}\sin\theta + 9a_{u\phi}B_u - 12a_\phi B_u^2 + 3a_\phi B_{uu} \\
&+ 12b_\theta B_u^2\sin\theta - 9b_{u\theta}B_u\sin\theta - 3b_\theta B_{uu}\sin\theta + 4e^{4B}B_\theta B_{u\theta} \\
&- e^{4B}B_{u\theta}\cot\theta + e^{4B}B_{u\theta\theta} - 4e^{4B}B_{u\phi}B_\phi\csc^2\theta - e^{4B}B_{u\phi\phi}\csc^2\theta\Big) \\
&+ ie^{-6B}\Big(-\frac{3}{2}a_{uu\phi}\sin\theta - \frac{3}{2}b_{uu\phi} - 12a_\theta B_u^2\sin\theta + 9a_{u\theta}B_u\sin\theta \\
&+ 3a_\theta B_{uu}\sin\theta + 9b_{u\phi}B_u - 12b_\phi B_u^2 + 3b_\phi B_{uu} - 4e^{4B}B_{u\theta}B_\phi\csc\theta \\
&- 2e^{4B}B_{u\phi}\csc\theta - 4e^{4B}B_\theta B_{u\phi}\csc\theta + 2e^{4B}B_{u\phi}\cot\theta\csc\theta\Big), \\
(\Psi_3^2)^0 =& \frac{e^{-4B}}{2\sqrt{2}}\Big(B_\theta(-24a_\phi B_u + 12a_{u\phi} + 24b_\theta B_u\sin\theta - 12b_{u\theta}\sin\theta \\
&- 2e^{4B}\csc^2\theta + 8e^{4B}B_{\theta\theta} + 4e^{4B}) + 24a_\theta B_u B_\phi + 12a_\phi B_u\cot\theta \\
&+ 6a_\phi B_{u\theta} - 12a_{u\theta}B_\phi - 6a_\theta B_{u\phi} - 6a_{u\phi}\cot\theta - 6b_{\theta\theta}B_u\sin\theta \\
&+ 24b_\phi B_u B_\phi\csc\theta - 6b_{\phi\phi}B_u\csc\theta + 6bB_u\sin\theta - 18b_\theta B_u\cos\theta \\
&- 6bB_u\csc\theta + 6bB_u\cos\theta\cot\theta - 6b_\theta B_{u\theta}\sin\theta - 6b_\phi B_{u\phi}\csc\theta \\
&- 12b_{u\phi}B_\phi\csc\theta - 3b_u\sin\theta + 3b_u\csc\theta - 3b_u\cos\theta\cot\theta + 9b_{u\theta}\cos\theta \\
&+ 3b_{u\theta\theta}\sin\theta + 3b_{u\phi\phi}\csc\theta + 2e^{4B}B_{\theta\theta}\cot\theta + 2e^{4B}B_{\theta\phi\phi}\csc^2\theta \\
&- 4e^{4B}B_{\phi\phi}\csc^2\theta(\cot\theta - 2B_\theta) + 8e^{4B}B_\theta^2\cot\theta + 2e^{4B}B_{\theta\theta\theta}\Big) \\
&+ \frac{ie^{-4B}}{2\sqrt{2}}\Big(-4B_\phi(6a_\phi B_u\csc\theta - 3a_{u\phi}\csc\theta - 6b_\theta B_u + 3b_{u\theta} \\
&+ 2e^{4B}B_{\theta\theta}\csc\theta + 2e^{4B}B_{\phi\phi}\csc^3\theta + e^{4B}\csc\theta + 2e^{4B}B_\theta\cot\theta\csc\theta) \\
&+ 6a_{\theta\theta}B_u\sin\theta + 6a_{\phi\phi}B_u\csc\theta - 6aB_u\sin\theta - 24a_\theta B_\theta B_u\sin\theta \\
&+ 18a_\theta B_u\cos\theta + 6aB_u\csc\theta - 6aB_u\cos\theta\cot\theta + 6a_\theta B_{u\theta}\sin\theta \\
&+ 12a_{u\theta}B_\theta\sin\theta + 6a_\phi B_{u\phi}\csc\theta + 3a_u\sin\theta - 3a_u\csc\theta \\
&+ 3a_u\cos\theta\cot\theta - 9a_{u\theta}\cos\theta - 3a_{u\theta\theta}\sin\theta - 3a_{u\phi\phi}\csc\theta - 24b_\phi B_\theta B_u \\
&+ 12b_\phi B_u\cot\theta + 6b_\phi B_{u\theta} - 6b_\theta B_{u\phi} + 12b_{u\phi}B_\theta - 6b_{u\phi}\cot\theta \\
&- 2e^{4B}B_{\theta\theta\phi}\csc\theta - 2e^{4B}B_{\theta\phi}\cot\theta\csc\theta - 2e^{4B}B_{\phi\phi\phi}\csc^3\theta\Big), \\
(\Psi_2^3)^0 =& \frac{e^{-6B}}{4}\Big(4e^{4B}M + 3(12a_\phi b_\theta B_u\sin\theta - 12a_\theta b_\phi B_u\sin\theta - 3a_\phi b_{u\theta}\sin\theta \\
&+ 3a_{u\theta}\sin\theta(a_\theta\sin\theta + b_\phi) + 3a_{u\phi}(a_\phi - b_\theta\sin\theta) + 3a_\theta b_{u\phi}\sin\theta \\
&- 4e^{4B}a_{\phi\phi}B_\phi\csc^2\theta - 8e^{4B}a_\phi B_\theta^2 + 16e^{4B}a_\theta B_\theta B_\phi + 8e^{4B}a_\phi B_\theta\cot\theta \\
&- 12e^{4B}a_\theta B_\phi\cot\theta + 8e^{4B}a_\phi B_\phi^2\csc^2\theta - 4e^{4B}a_{\theta\theta}B_\phi - 6a_\theta^2 B_u\sin^2\theta \\
&- 6a_\phi^2 B_u - 4e^{4B}b_{\theta\theta}B_\theta\sin\theta - 8e^{4B}b_\theta B_\phi^2\csc\theta + 16e^{4B}b_\phi B_\theta B_\phi\csc\theta
\end{aligned}
$$

$$
\begin{aligned}
&- 8e^{4B}b_\phi B_\phi \cot\theta \csc\theta - 4e^{4B}b_{\phi\phi}B_\theta \csc\theta + 8e^{4B}b_\theta B_\theta^2 \sin\theta \\
&+ 4be^{4B}B_\theta \sin\theta - 12e^{4B}b_\theta B_\theta \cos\theta + 4be^{4B}B_\theta \cos\theta \cot\theta \\
&- 4be^{4B}B_\theta \csc\theta - 6b_\theta^2 B_u \sin^2\theta - 6b_\phi^2 B_u + 3b_\theta b_{u\theta}\sin^2\theta + 3b_{u\phi}b_\phi)\Big) \\
&+ \frac{3ie^{-6B}}{4}\Big(a_\theta \sin\theta(-3a_{u\phi} + 3b_{u\theta}\sin\theta + e^{4B}(-4B_{\phi\phi}\csc^2\theta + 8B_\theta \cot\theta \\
&+ 4B_{\theta\theta} - 3\cot^2\theta + 3)) - 3a_{u\theta}b_\theta \sin^2\theta + 3a_\phi b_{u\phi} - 3a_{u\phi}b_\phi \\
&+ e^{4B}a_{\theta\theta}(4B_\theta \sin\theta - 5\cos\theta) - e^{4B}a_{\theta\theta\theta}\sin\theta + 8e^{4B}a_\phi B_{\theta\phi}\csc\theta \\
&- e^{4B}a_{\theta\phi\phi}\csc\theta + 4e^{4B}a_{\phi\phi}B_\theta \csc\theta - 2e^{4B}a_{\phi\phi}\cot\theta\csc\theta \\
&- 4ae^{4B}B_\theta \sin\theta + 3ae^{4B}\cos\theta + 4ae^{4B}B_\theta \csc\theta + 3ae^{4B}\cos\theta\cot^2\theta \\
&- 4ae^{4B}B_\theta \cos\theta\cot\theta - 3ae^{4B}\cot\theta\csc\theta + 3a_{u\theta}a_\phi \sin\theta - 8e^{4B}b_\theta B_{\theta\phi} \\
&+ e^{4B}b_{\theta\phi}\cot\theta - 4e^{4B}b_{\phi\phi}B_\phi \csc^2\theta - 4e^{4B}b_\phi B_{\phi\phi}\csc^2\theta - 3e^{4B}b_\phi \cot^2\theta \\
&+ 4be^{4B}B_\phi \cot^2\theta + 4e^{4B}b_\phi B_\theta \cot\theta - 4e^{4B}b_\theta B_\phi \cot\theta + 3e^{4B}b_\phi \csc^2\theta \\
&- 4be^{4B}B_\phi \csc^2\theta + e^{4B}b_{\phi\phi\phi}\csc^2\theta + 4e^{4B}b_\phi B_{\theta\theta} - 4e^{4B}b_{\theta\theta}B_\phi \\
&+ e^{4B}b_{\theta\phi\phi} - e^{4B}b_\phi + 4be^{4B}B_\phi + 3b_{u\theta}b_\phi \sin\theta - 3b_\theta b_{u\phi}\sin\theta\Big),
\end{aligned}
$$

$$
\begin{aligned}
(\Psi_1^4)^0 =& \frac{3e^{-4B}\csc\theta}{32\sqrt{2}}\Big(-16e^{2B}\sin\theta U_3 + 36a_\theta \sin\theta(3a_{\theta\theta}\sin^2\theta + 2a_{\phi\phi} \\
&- 10b_\phi B_\theta \sin\theta + b_{\theta\phi}\sin\theta + 11b_\phi\cos\theta) + 180a_\phi b_\theta B_\theta \\
&- 180a_\phi b_\theta B_\theta \cos 2\theta + 54a_\phi b_{\theta\theta}\cos 2\theta - 54a_{\theta\theta}b_\phi \cos 2\theta \\
&- 36a_{\theta\phi}\sin\theta(b_\theta \sin\theta - a_\phi) - 198a_\phi b_\theta \sin 2\theta - 54a_\phi b_{\theta\theta} \\
&+ 54a_{\theta\theta}b_\phi + 72a_{\phi\phi}b_\phi - 72a_\phi b_{\phi\phi} - 180a_\phi^2 B_\theta \sin\theta + 144a_\phi^2\cos\theta \\
&+ 144b_\phi^2\cos\theta - 36a_\theta^2\sin^2\theta(5B_\theta\sin\theta - 7\cos\theta) - 180b_\phi^2 B_\theta\sin\theta \\
&- 135b_\theta^2 B_\theta \sin\theta + 45b_\theta^2 B_\theta \sin 2\theta + 81b_\theta b_{\theta\theta}\sin\theta - 27b_\theta b_{\theta\theta}\sin 2\theta \\
&+ 36b_{\theta\phi}b_\phi \sin\theta + 72b_\theta b_{\phi\phi}\sin\theta + 63b_\theta^2\cos\theta - 63b_\theta^2\cos 2\theta\Big) \\
&\frac{3ie^{-4B}\csc\theta}{16\sqrt{2}}\Big(-8e^{2B}\sin\theta W_3 + 18a_\theta\sin\theta(a_{\theta\phi}\sin\theta + 2a_\phi\cos\theta \\
&- 10b_\phi B_\phi - b_{\theta\theta}\cos 2\theta + b_{\theta\theta} + 3b_{\phi\phi}) + 180a_\phi b_\theta B_\phi \sin\theta \\
&+ 36a_{\theta\theta}\sin^2\theta(a_\phi - b_\theta\sin\theta) - 18a_\phi b_{\theta\phi}\sin\theta + 18a_{\theta\phi}b_\phi\sin\theta \\
&- 54a_{\phi\phi}b_\theta\sin\theta - 90a_\theta^2 B_\phi\sin^2\theta - 90a_\phi^2 B_\phi + 54a_\phi a_{\phi\phi} - 45b_\theta^2 B_\phi \\
&+ 45b_\theta^2 B_\phi\cos 2\theta - 90b_\phi^2 B_\phi - 18b_{\theta\theta}b_\phi\cos 2\theta + 9b_\theta b_{\theta\phi} \\
&- 9b_\theta b_{\theta\phi}\cos 2\theta + 18b_\theta b_\phi \sin 2\theta + 18b_{\theta\theta}b_\phi + 54b_\phi b_{\phi\phi}\Big),
\end{aligned}
$$

$$
(\Psi_0^5)^0 = 6(C + iD), \quad (\Psi_0^5)^1 = 0.
$$

Acknowledgement. This work is partially supported by the National Science Foundation of China (grant 11571345) and the project of mathematics and interdisciplinary sciences of Chinese Academy of Sciences.

References

[1] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, Waves from axi-symmetric isolated systems, Proc. R. Soc. London A **269**, 21 (1962).
[2] R. K. Sachs, Waves in asymptotically flat space-time, Proc. R. Soc. London A **270**, 103 (1962).
[3] R. Penrose, Zero rest mass fields including gravitation: asymptotic behavior, Proc. R. Soc. A **284**, 159 (1965).
[4] W-L. Huang, S.T. Yau, X. Zhang, Positivity of the Bondi mass in Bondi's radiating space-times. Rend. Lincei Mat. Appl. **17**, 335 (2006).
[5] X. Zhang, Recent progress on the positive energy theorem, Int. J. Mod. Phys. A **30**, 1545018 (2015).
[6] H. Ge, M. Luo, Q. Su, D. Wang, X. Zhang, Bondi-Sachs metrics and photon rockets, Gen. Relativ. Gravit. **43**, 2729 (2011).
[7] X. He, Z. Cao, New Bondi-type outgoing boundary condition for the Einstein equations with cosmological constant, Int. J. Mod. Phys. D **24**, 1550081 (2015).
[8] A. Ashtekar, B. Bonga, A. Kesavan, Asymptotics with a positive cosmological constant: I. Basic framework, Class. Quantum Grav. **32**, 025004 (2015).
[9] A. Ashtekar, B. Bonga, A. Kesavan, Asymptotics with a positive cosmological constant. II. Linear fields on de Sitter spacetime, Phys. Rev. D **92**, 044011 (2015).
[10] A. Ashtekar, B. Bonga, A. Kesavan, Asymptotics with a positive cosmological constant. III. The quadrupole formula, Phys. Rev. D **92**, 104032 (2015).
[11] A. Ashtekar, B. Bonga, A. Kesavan, Gravitational Waves from Isolated Systems: Surprising Consequences of a Positive Cosmological Constant, Phys. Rev. Lett. **116**, 051101 (2016).
[12] N. T. Bishop, Gravitational waves in a de Sitter universe, Phys. Rev. D **93**, 044025 (2016).
[13] G. Date, S. J. Hoque, Gravitational Waves from Compact Sources in de Sitter Background, Phys. Rev. D **94**,064039 (2016).
[14] G. Date, S. J. Hoque, Cosmological Horizon and the Quadrupole Formula in de Sitter Background, arXiv:1612.09511v2 [gr-qc] 5 Apr 2017.

Institute of Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

E-mail address: fangquanxie@amss.ac.cn, xzhang@amss.ac.cn